
Main Contraindications for Use

General Contraindications

1. **Pregnancy and breastfeeding** – lack of studies confirming the safety of use in pregnant and breastfeeding women.
2. **Autoimmune diseases** – e.g., systemic lupus erythematosus, scleroderma, due to the possibility of unpredictable immune system responses.
3. **Allergy to the product's ingredients** – hypersensitivity to poly-L-lactic acid or other components of the preparation.
4. **Infections** – local skin infections (e.g., herpes, cystic acne) or systemic infections (e.g., flu, viral infections).
5. **Cancer** – both active and recently treated conditions.
6. **Skin inflammation** – at the intended injection site.

Local Contraindications

1. **Skin damage** – e.g., wounds, ulcers, hypertrophic scars at the planned treatment area.
2. **Tendency to scar formation** – particularly hypertrophic scarring or keloids.
3. **Active skin diseases** – e.g., psoriasis, eczema, active acne in the treatment area.
4. **Presence of implants in the treatment area** – e.g., silicone implants or other foreign bodies at the planned injection site.

Procedural Contraindications

1. **Application in contraindicated areas** – e.g., sensitive regions such as eyelids or vermillion border.
2. **Combined therapies** – the procedure should not be performed simultaneously with other invasive treatments in the same area (e.g., HA fillers, lasers, microneedling) if recovery time is insufficient.

Recommendations

Before proceeding with a procedure using AesPLLa 365 mg, a thorough medical history and skin evaluation must be conducted.

If there are any uncertainties regarding contraindications, consultation with a medical specialist is strongly advised.