
Contraindications

- Severe allergies with a history of severe reactions (anaphylaxis). Use of Restylane Refyne may cause an allergic reaction.
- Allergy to the anesthetic lidocaine or any of the proteins used to make hyaluronic acid; the acid in Restylane Refyne (bacterial proteins). Precautions
- Restylane Refyne helps treat facial wrinkles - natural wrinkles and folds on the face that can occur due to aging. Restylane Refyne has not been tested to treat other types of wrinkles.
- If you experience vision changes, stroke symptoms (including sudden difficulty speaking, numbness or weakness in the face, arms or legs, difficulty walking, facial drooping, severe headache, dizziness or confusion), a white appearance of the skin or unusual pain during or shortly after treatment, notify your doctor immediately.
- Use of Restylane Refyne if you have skin ulcers, pimples, rashes, hives, cysts or infections should be postponed until it has fully healed.
- Using Restylane Refyne in such cases may delay healing or worsen skin problems.
- Restylane Refyne is intended for adults 22 years of age and older.
- The safety of Restylane Refyne has not been studied in women who are pregnant or breastfeeding.
- Talk to your doctor before using Restylane Refyne if you are pregnant and planning to become pregnant or breastfeeding
- If you are using Restylane Refyne while taking any medications that reduce the body's natural defense system ("immunosuppressants")
- Some medications, such as aspirin and warfarin, can thin the blood, which can increase the risk of: bleeding or prolong any bleeding that occurs. These drugs also increase the risk of bruising or bleeding at the injection site. Tell your doctor about any medications you are taking.
- As with any injection procedure, you may have a higher risk of bruising or bleeding at the injection site if you have a hemorrhagic diathesis.
- If you have any history of scarring, especially thick and stiff scars, or any skin color (pigmentation), talk to your doctor about it before the procedure. scarring and skin color changes can occur with hyaluronic acid fillers in general.
- When Restylane Refyne was tested in patients with a variety of skin tones (pale to dark), there were no reports from patients with excessive scarring (keloids) or skin color changes. ☒ The use of Restylane Refyne in skin therapies, such as laser treatments or chemical peels, can: lead to other side effects, such as redness, swelling, warmth and/or pain in the skin (inflammatory reaction). Inform your doctor if you have had any of these procedures.